

11. Problemset “Quantum Algebra & Dynamics”

January 8, 2016

Fock Representation(s) revisited

11.1 Driven Oscillators

Consider this caricature

$$H = \sum_i \omega_i a_i^* a_i + g \sum_i (\bar{j}_i a_i + j_i a_i^*) \quad (1)$$

of a scalar field ϕ coupled to a classical source j

$$H = \int dx \left(\frac{1}{2} \pi^2(x) + \frac{1}{2} (\nabla \phi)^2(x) + \frac{m^2}{2} \phi^2(x) + j(x) \phi(x) \right).$$

1. Find annihilation and creation operators A_i , A_i^* and energies E_i as functions of a_i , a_i^* , ω_i , j_i and \bar{j}_i , such that (1) can be written

$$H = \sum_i E_i A_i^* A_i + \text{const.} \quad (2)$$

Determine the shift in the ground state energy.

2. Find the unitary intertwiners U with

$$A_i = U a_i U^* \quad (3)$$

as a function of ω_i , j_i and \bar{j}_i .

3. Under which conditions on ω_i , j_i and \bar{j}_i lies the interacting ground state Ω with

$$A_i \Omega = 0 \quad (4)$$

in the Fock representation defined by the a_i and a_i^* ? Compute Ω in this case.

4. Under which conditions on ω_i , j_i and \bar{j}_i are the U well defined unitary operators in the Fock representation?

11.2 Normal Ordering the Squeeze Operator

Prove that the unitary creation operators

$$U_s(\zeta) = e^{\frac{1}{2}(\bar{\zeta}aa - \zeta a^*a^*)} \quad (5)$$

for squeezed states can be normal ordered as

$$U_s(\zeta) = e^{-\frac{\zeta}{2} \frac{\tanh|\zeta|}{|\zeta|} a^*a^*} e^{-\ln(\cosh|\zeta|)(a^*a + \frac{1}{2})} e^{\frac{\bar{\zeta}}{2} \frac{\tanh|\zeta|}{|\zeta|} aa}. \quad (6)$$

For convenience, introduce

$$A = \frac{1}{2}aa \quad (7a)$$

$$A^* = \frac{1}{2}a^*a^* \quad (7b)$$

$$N = a^*a + \frac{1}{2} \quad (7c)$$

to write

$$U_s(\zeta) = e^{\bar{\zeta}A - \zeta A^*} = e^{-\zeta \frac{\tanh|\zeta|}{|\zeta|} A^*} e^{-\ln(\cosh|\zeta|)N} e^{\bar{\zeta} \frac{\tanh|\zeta|}{|\zeta|} A}. \quad (8)$$

If the Lie algebra of the exponents in (14) closes, the Baker-Campbell-Hausdorff formula shows that we can express the exponents as linear combinations of the generators. Then it suffices to compute the coefficients in a matrix representation.

1. Show that A , A^* and N form a closed Lie algebra.
2. Show that π with

$$\pi(A) = \begin{pmatrix} 0 & 0 \\ -1 & 0 \end{pmatrix} \quad (9a)$$

$$\pi(A^*) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} \quad (9b)$$

$$\pi(N) = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} \quad (9c)$$

is a representation of the Lie algebra. (NB: $\pi(A^*) \neq (\pi(A))^\dagger$, but we will never use that in the following!)

3. Compute the matrix exponentials in (14) to prove the normal ordering formula.