

5. Problemset “Quantum Algebra & Dynamics”

November 16, 2018

States

NB: the notions of *positive linear functional* and *state* will be introduced in the lecture on Wednesday, November 21, 2015.

The space of continuous linear functionals $\omega : \mathcal{A} \rightarrow \mathbf{C}$ on the C^* -algebra \mathcal{A} is denoted \mathcal{A}^* .

We can define a natural norm on \mathcal{A}^* by

$$\|\omega\| = \sup_{A \in \mathcal{A}, \|A\|=1} |\omega(A)|. \quad (1)$$

A linear functional $\omega : \mathcal{A} \rightarrow \mathbf{C}$ on the C^* -algebra \mathcal{A} is called positive, iff

$$\forall A \in \mathcal{A} : \omega(A^*A) \geq 0. \quad (2)$$

A positive $\omega : \mathcal{A} \rightarrow \mathbf{C}$ with $\|\omega\| = 1$ is called a state.

5.1 Spins

Consider again the C^* -algebra \mathcal{M}_2 of 2×2 -Matrices parametrized by four complex numbers (a_0, \vec{a}) , using the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}. \quad (3)$$

Define a family of linear functionals

$$\begin{aligned} \omega_{a_0, \vec{a}} : \mathcal{M}_2 &\rightarrow \mathbf{C} \\ M(b_0, \vec{b}) &\mapsto \text{tr}(M(b_0, \vec{b})\rho(a_0, \vec{a})) \end{aligned} \quad (4)$$

for suitable $\rho(a_0, \vec{a}) \in \mathcal{M}_2$. Derive the conditions on (a_0, \vec{a}) for $\omega_{a_0, \vec{a}}$ to be ...

1. ... continuous?
2. ... positive?
3. ... a state?
4. ... maximal in the sense that $\omega_{a_0, \vec{a}}$ can *not* be written

$$\omega_{a_0, \vec{a}} = p\omega_{b_0, \vec{b}} + (1-p)\omega_{c_0, \vec{c}} \quad (5)$$

with $0 < p < 1$ and $\omega_{b_0, \vec{b}}$ and $\omega_{c_0, \vec{c}}$ states?

5.2 Circle

Consider the algebra $C(S^1)$ of bounded complex valued continuous functions $f : S^1 \rightarrow \mathbf{C}$ on the unit circle.

1. Show that $\|f\| = \sup |f(x)|$ turns $C(S^1)$ into a C^* -algebra.
2. Define linear functionals $\omega : C(S^1) \rightarrow \mathbf{C}$ via

$$\omega(f) = \int_0^{2\pi} \frac{d\phi}{2\pi} \overline{\omega(\phi)} f(\phi). \quad (6)$$

What are the conditions on $\omega : S^1 \rightarrow \mathbf{C}$ for ω to be ...

- (a) ... continuous?
- (b) ... positive?
- (c) ... a state?
- (d) ... maximal in the sense that ω can *not* be written

$$\omega = p\omega_1 + (1-p)\omega_2 \quad (7)$$

with $0 < p < 1$ and $\omega_{1/2}$ states as in (6)?

3. Are there states $\omega : C(S^1) \rightarrow \mathbf{C}$ that can not be written as in (6)?
4. If yes, give examples and repeat the second subproblem!