## 7. Problemset "Quantum Algebra & Dynamics"

November 30, 2018

## **GNS** Construction

## 7.1 Spins

Consider again the  $C^*$ -algebra  $\mathcal{M}_2$  of  $2 \times 2$ -Matrices  $M(a_0, \vec{a})$  parametrized by four complex numbers  $(a_0, \vec{a})$ , using the Pauli matrices

$$\sigma_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \ \sigma_2 = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \ \sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$
 (1)

with

$$M(a_0, \vec{a}) = a_0 \mathbf{1} + \vec{a} \vec{\sigma} \,. \tag{2}$$

As you have shown in problem 5.1, the states on  $\mathcal{M}_2$  can be parametrized by three real numbers  $\vec{\alpha}$  with  $|\vec{\alpha}| \leq 1$  and

$$\omega_{\vec{\alpha}} : \mathcal{M}_2 \to \mathbf{C}$$

$$M \mapsto \frac{1}{2} \operatorname{tr}(M\rho(1, \vec{\alpha})) \tag{3}$$

using  $\rho(\alpha_0, \vec{\alpha}) \in \mathcal{M}_2$ . The pure states are those with  $|\vec{\alpha}| = 1$ .

1. Perform the GNS construction of cyclic representations  $(\mathcal{H}_{\vec{\alpha}}, \pi_{\vec{\alpha}}, \Omega_{\vec{\alpha}}) = (\mathcal{H}_{\omega_{\vec{\alpha}}}, \pi_{\omega_{\vec{\alpha}}}, \Omega_{\omega_{\vec{\alpha}}})$  in the special cases

$$\vec{\alpha} = (0, 0, 0) \tag{4a}$$

$$\vec{\alpha} = (0, 0, a) \qquad \text{(with } |a| < 1) \tag{4b}$$

$$\vec{\alpha} = (0, 0, 1)$$
 . (4c)

2. Give the concrete matrix realizations of

$$\pi_{\vec{\alpha}}(M(a_0, \vec{a})). \tag{5}$$

- 3. Show that (4c) leads to an irreducible and (4a) and (4b) to reducible representations. Determine the invariant subspace(s) in the latter cases.
- 4. Try to repeat the above problems in the general case

$$|\vec{\alpha}| < 1 \tag{6a}$$

$$|\vec{\alpha}| = 1. \tag{6b}$$